

CHAPTER 5

Checklist of meso- and macrozooplankton species of the Baltic Sea

Table 5.1: Meso- and macrozooplankton species in the Baltic Sea: **BP** – Baltic Proper; **WBS** – Western Baltic Sea; **NBS** – Northern Baltic Sea, **SBS** – Southern Baltic Sea; **EBS** – Eastern Baltic Sea (“+” present; no sign = species not found). After Telesh et al. (2009), with additions and modifications.

No	Taxa	BP ¹	WBS ²	NBS ³	SBS ⁴	EBS ⁵
Cnidaria, Anthomedusae						
1	<i>Euphysa aurata</i> Forbes, 1848	+	+			
2	<i>Euphysa tentaculata</i> Linko, 1905	+	+			
3	<i>Halitholus cirratus</i> Hartlaub, 1913	+	+		+	
4	<i>Hybocodon prolifer</i> L. Agassiz, 1862	+	+			
5	<i>Rathkea octopunctata</i> (M. Sars, 1835)	+	+			
6	<i>Sarsia tubulosa</i> (M. Sars, 1835)	+	+			
Cnidaria, Trachymedusae						
7	<i>Aglantha digitalis</i> (O.F. Müller, 1776)	+	+			
Cnidaria, Scyphomedusae						
8	<i>Aurelia aurita</i> (Linnaeus, 1758)	+	+	+	+	
9	<i>Clytia hemisphaerica</i> (Linnaeus, 1767)	+				
10	<i>Cyanea capillata</i> (Linnaeus, 1758)		+			
11	<i>Haliclystus auricula</i> O. Fabricius, 1780		+			
12	<i>Lucernaria quadricornis</i> O.F. Müller, 1776		+			
13	<i>Rhizostoma octopus</i> (Mayer, 1910)		+			
Cnidaria, Leptomedusae						
14	<i>Melicertum octocostatum</i> (M. Sars, 1835)	+	+		+	
15	<i>Obelia geniculata</i> (Linnaeus, 1758)	+	+		+	

No	Taxa	BP ¹	WBS ²	NBS ³	SBS ⁴	EBS ⁵
Ctenophora						
16	<i>Beroe cucumis</i> Fabricius, 1780	+	+			
17	<i>Beroe gracilis</i> Künne, 1939	+	+			
18	<i>Bolinopsis infundibulum</i> O.F. Müller, 1776	+	+			
19	<i>Mnemiopsis leidyi</i> A. Agassiz, 1865	+	+	+	+	
20	<i>Pleurobrachia pileus</i> (O.F. Müller, 1776)	+	+	+	+	+
Turbellaria						
21	<i>Alaurina composita</i> Metschnikow, 1865		+			
Rotifera						
22	<i>Anuraeopsis fissa</i> (Gosse, 1851)			+		
23	<i>Asplanchna</i> sp.			+		
24	<i>Asplanchna brightwelli</i> Gosse, 1850		+			+
25	<i>Asplanchna priodonta</i> Gosse, 1850		+		+	+
26	<i>Asplanchna seiboldi</i> (Leydig, 1854)		+			
27	<i>Bdelloidea</i> indet.			+		+
28	<i>Brachionus</i> sp.			+	+	+
29	<i>Brachionus angularis</i> Gosse, 1851		+	+	+	+
30	<i>Brachionus calyciflorus</i> Pallas, 1766		+	+	+	+
31	<i>Brachionus plicatilis</i> O.F. Müller, 1786		+			
32	<i>Brachionus quadridentatus</i> Hermann, 1783		+	+		
33	<i>Brachionus rubens</i> Ehrenberg, 1838		+			
34	<i>Brachionus urceus</i> (Linnaeus, 1758)		+	+		
35	<i>Cephalodella catellina</i> (O.F. Müller, 1786))		+			
36	<i>Cephalodella megalcephala</i> (Glasscott, 1893)		+			
37	<i>Collotheca</i> sp.					+
38	<i>Collotheca mutabilis</i> (Hudson, 1885)		+			
39	<i>Collotheca ornata</i> (Ehrenberg, 1832)		+			
40	<i>Collotheca pelagica</i> (Rousselet, 1893)					+
41	<i>Colurella</i> sp.		+	+		+
42	<i>Conochilus unicornis</i> Rousselet, 1892				+	
43	<i>Dicranophorus</i> sp.				+	
44	<i>Encentrum pachypus</i> Remane, 1949		+			+
45	<i>Euchlanis</i> sp.					+
46	<i>Euchlanis dilatata</i> Ehrenberg, 1832		+	+	+	+
47	<i>Filinia brachiata</i> (Rousselet, 1901)			+		

No	Taxa	BP ¹	WBS ²	NBS ³	SBS ⁴	EBS ⁵
48	<i>Filinia longiseta</i> (Ehrenberg, 1834)		+	+	+	+
49	<i>Filinia terminalis</i> (Plate, 1886)		+	+	+	
50	<i>Hexarthra fennica</i> (Levander, 1892)		+			+
51	<i>Kellicottia longispina</i> (Kellicott, 1879)			+	+	+
52	<i>Keratella cochlearis</i> (Gosse, 1851)	+	+	+	+	+
53	<i>Keratella cochlearis baltica</i> (Sokolova, 1927)	+	+	+	+	+
54	<i>Keratella cochlearis recurispina</i> (Jägerskiöld, 1894)	+		+		+
55	<i>Keratella cochlearis tecta</i> (Gosse, 1851)		+	+	+	+
56	<i>Keratella cruciformis eichwaldi</i> (Levander, 1894)	+	+	+	+	+
57	<i>Keratella quadrata</i> (O.F. Müller, 1786)	+	+	+	+	+
58	<i>Keratella quadrata platei</i> (Jägerskiöld, 1894)	+		+		+
59	<i>Lecane</i> sp.			+		+
60	<i>Lecane lamellata</i> (Daday, 1893)		+			
61	<i>Lecane luna</i> (O.F. Müller, 1776)					+
62	<i>Lecane lunaris</i> (Ehrenberg, 1832)		+			
63	<i>Lepadella</i> sp.			+		
64	<i>Monommata</i> sp.			+		
65	<i>Mytilina mucronata</i> (O.F. Müller, 1773)		+			
66	<i>Notholca</i> sp.				+	+
67	<i>Notholca caudata</i> Carlin, 1943			+		
68	<i>Notholca acuminata</i> (Ehrenberg, 1832)	+	+			+
69	<i>Notholca acuminata extensa</i> Oloffsson, 1918		+			
70	<i>Notholca acuminata marina</i> Focke, 1961		+			
71	<i>Notholca labis</i> Gosse, 1887					+
72	<i>Notholca squamula</i> (O.F. Müller, 1786)					+
73	<i>Notholca squamula salina</i> Focke, 1961		+			
74	<i>Notholca striata</i> (O.F. Müller, 1786)	+	+	+		
75	<i>Philodina</i> sp.			+		
76	<i>Ploesoma truncatum</i> (Levander, 1894)			+		+
77	<i>Polyarthra</i> sp.				+	+
78	<i>Polyarthra dolichoptera</i> Idelson, 1925	+	+	+	+	+
79	<i>Polyarthra major</i> Burckhardt, 1900			+		
80	<i>Polyarthra remata</i> Skorikov, 1896			+		
81	<i>Polyarthra vulgaris</i> Carlin, 1943		+	+		+
82	<i>Pompholyx sulcata</i> Hudson, 1885			+		
83	<i>Proales</i> sp.		+			+

No	Taxa	BP ¹	WBS ²	NBS ³	SBS ⁴	EBS ⁵
84	<i>Proales reinhardti</i> (Ehrenberg, 1834)			+		
85	<i>Synchaeta</i> sp.	+		+	+	+
86	<i>Synchaeta baltica</i> Ehrenberg, 1834	+	+	+	+	+
87	<i>Synchaeta cecilia</i> Rousselet, 1902		+	+		
88	<i>Synchaeta curvata</i> Lie-Pettersen, 1905	+	+	+		+
89	<i>Synchaeta fennica</i> Rousselet, 1909	+	+		+	+
90	<i>Synchaeta grimpei</i> Remane, 1929		+			
91	<i>Synchaeta gyrina</i> Hood, 1887	+				+
92	<i>Synchaeta littoralis</i> Rousselet, 1902		+	+		+
93	<i>Synchaeta monopus</i> Plate, 1889	+	+	+	+	+
94	<i>Synchaeta pectinata</i> Ehrenberg, 1832		+			
95	<i>Synchaeta triophthalma</i> Lauterborn, 1894	+	+			
96	<i>Synchaeta vorax</i> Rousselet, 1902					+
97	<i>Testudinella clypeata</i> (O.F. Müller, 1786)		+			+
98	<i>Trichocerca</i> sp.		+	+	+	
99	<i>Trichocerca capucina</i> (Wierzejski et Zacharias, 1893)			+		+
100	<i>Trichocerca dixon-nutalli</i> Jennings, 1903		+			
101	<i>Trichocerca marina</i> (Daday, 1890)		+	+		+
102	<i>Trichocerca pusilla</i> (Lauterborn, 1898)					+
103	<i>Trichocerca (Diurella) similis</i> (Wierzejski, 1893)			+		
104	<i>Trichotria pocillum</i> (O.F. Müller, 1776)					+
Cladocera						
105	<i>Acroperus harpae</i> (Baird, 1834) °				+	+
106	<i>Alona</i> sp.			+	+	
107	<i>Alona affinis</i> Leydig, 1860 °				+	+
108	<i>Alona costata</i> G.O. Sars, 1862 °				+	+
109	<i>Alona guttata</i> G.O. Sars, 1862 °				+	+
110	<i>Alona intermedia</i> G.O. Sars, 1862		+			+
111	<i>Alona quadrangularis</i> (O.F. Müller, 1776)				+	+
112	<i>Alona rectangula</i> Sars, 1861		+		+	+
113	<i>Alonella excisa</i> (Fischer, 1854) °				+	+
114	<i>Alonella exigua</i> (Lilljeborg, 1853) °				+	
115	<i>Alonella nana</i> (Baird, 1850) °					+
116	<i>Alonopsis elongata</i> Sars, 1862				+	+
117	<i>Bosmina crassicornis</i> P.E. Müller, 1867					+
118	<i>Bosmina longirostris</i> (O.F. Müller, 1776)	+	+	+	+	+

No	Taxa	BP ¹	WBS ²	NBS ³	SBS ⁴	EBS ⁵
119	<i>Bythotrephes</i> sp.					+
120	<i>Camptocercus rectirostris</i> Schoedler, 1862 °				+	+
121	<i>Camptocercus lilljeborgii</i> Schoedler **, °					+
122	<i>Cercopagis pengoi</i> (Ostroumov, 1891)	+		+	+	+
123	<i>Ceriodaphnia</i> sp.			+	+	+
124	<i>Ceriodaphnia laticaudata</i> P.E. Müller, 1867		+			
125	<i>Ceriodaphnia pulchella</i> G.O. Sars, 1862				+	+
126	<i>Ceriodaphnia quadrangula</i> (O.F. Müller, 1785)				+	+
127	<i>Ceriodaphnia reticulata</i> (Jurine, 1820)		+		+	
128	<i>Ceriodaphnia setosa</i> Matile **, °				+	
129	<i>Chydorus latus</i> Sars, 1862 °					+
130	<i>Chydorus ovalis</i> Kurz, 1874 °				+	+
131	<i>Chydorus piger</i> G.O. Sars, 1862 °				+	
132	<i>Chydorus sphaericus</i> (O.F. Müller, 1785)		+	+	+	+
133	<i>Cornigerius maeoticus</i> Pengo, 1879					+
134	<i>Daphnia</i> sp.				+	+
135	<i>Daphnia cristata</i> G.O. Sars, 1861			+	+	+
136	<i>Daphnia cucullata</i> G.O. Sars, 1862	+		+	+	+
137	<i>Daphnia curvirostris</i> Eylman, 1887 °				+	
138	<i>Daphnia galeata</i> G.O. Sars, 1864		+			
139	<i>Daphnia longispina</i> (O.F. Müller, 1785)		+		+	+
140	<i>Daphnia magna</i> Straus, 1820		+			
141	<i>Daphnia pulex</i> Leydig, 1860 °				+	+
142	<i>Diaphanosoma brachyurum</i> (Liévin, 1848)		+	+	+	+
143	<i>Diaphanosoma mongolianum</i> Ueno, 1938		+			
144	<i>Disparalona rostrata</i> (Koch, 1841) °				+	+
145	<i>Eubosmina coregoni</i> Baird, 1857	+	+	+	+	+
146	<i>Eubosmina longispina</i> (Leidig, 1860) (Syn.*: <i>Bosmina longispina</i> Leidig, 1860; <i>Bosmina coregoni maritima</i> sensu Purasjoki, 1958)	+	+	+	+	+
147	<i>Eubosmina maritima</i> (P.E. Müller, 1867) (Syn.: <i>Bosmina maritima</i> P.E. Müller, 1867)	+	+	+	+	+
148	<i>Eurycercus lamellatus</i> (O.F. Müller, 1776)		+		+	+
149	<i>Evadne anonyx</i> G.O. Sars, 1897 [May be confused with <i>E. nordmanni</i>]				+	+
150	<i>Evadne nordmanni</i> Lovén, 1836	+	+	+	+	+
151	<i>Evadne spinifera</i> P.E. Müller, 1867		+			
152	<i>Graptoleberis testudinaria</i> (Fischer, 1848) °				+	+
153	<i>Holopedium gibberum</i> Zaddach, 1855 °					+

No	Taxa	BP ¹	WBS ²	NBS ³	SBS ⁴	EBS ⁵
154	<i>Ilyocryptus acutifrons</i> G. O. Sars, 1862 °				+	+
155	<i>Ilyocryptus agilis</i> Kurz, 1878 °					+
156	<i>Ilyocryptus sordidus</i> (Liévin, 1848) °					+
157	<i>Kurzia latissima</i> (Kurz, 1874) °					+
158	<i>Lathonura rectirostris</i> (O. F. Müller, 1776) °					+
159	<i>Latona setifera</i> (O. F. Müller, 1776) °				+	+
160	<i>Leptodora kindtii</i> (Focke, 1844)			+	+	+
161	<i>Leydigia leydigi</i> (Leydig, 1860) °				+	+
162	<i>Limnosida frontosa</i> Sars, ... **, °					+
163	<i>Macrothrix rosea</i> (Liévin, 1848) °					+
164	<i>Moina brachiata</i> (Jurine, 1820) °				+	+
165	<i>Monospilus dispar</i> G. O. Sars, 1861 °				+	+
166	<i>Ophryoxus gracilis</i> G. O. Sars, 1861 °					+
167	<i>Oxyurella tenuicaudis</i> (G. O. Sars, 1862) °				+	+
168	<i>Picripleuroxus laevis</i> (G. O. Sars, 1861) ° (Syn.: <i>Pleuroxus laevis</i> G. O. Sars, 1861)					+
169	<i>Picripleuroxus striatus</i> (Schödler, 1863) ° (Syn.: <i>Pleuroxus striatus</i> Schödler, 1863)					+
170	<i>Pleopsis polyphemoides</i> (Leuckart, 1859)	+	+	+	+	+
171	<i>Pleuroxus aduncus</i> (Jurine, 1820) °				+	+
172	<i>Pleuroxus trigonellus</i> (O. F. Müller, 1776) °				+	+
173	<i>Pleuroxus truncatus</i> (O. F. Müller, 1785) °				+	+
174	<i>Pleuroxus uncinatus</i> Baird, 1850 °				+	+
175	<i>Podon intermedius</i> Lilljeborg, 1853	+	+	+	+	+
176	<i>Podon leuckartii</i> (G.O. Sars, 1862)	+	+	+	+	+
177	<i>Polyphemus pediculus</i> (Linnaeus, 1761)					+
178	<i>Pseudochydorus globosus</i> (Baird, 1843) °				+	+
179	<i>Rhynchotalona falcata</i> (G. O. Sars, 1861) °				+	+
180	<i>Scapholeberis mucronata</i> (O. F. Müller, 1776) °				+	+
181	<i>Sida crystallina</i> (O.F. Müller, 1776)					+
182	<i>Simocephalus exspinosus</i> (DeGeer, 1778) °				+	+
183	<i>Simocephalus serrulatus</i> (Koch, 1841) °					+
184	<i>Simocephalus vetulus</i> (O. F. Müller, 1776) °				+	+
Copepoda, Calanoida						
185	<i>Acanthodiptomus denticornis</i> (Wierzejski, 1887) °					+
186	<i>Acartia bifilosa</i> (Giesbrecht, 1881)	+	+	+	+	+
187	<i>Acartia clausi</i> Giesbrecht, 1889	+	+	+	+	+

No	Taxa	BP ¹	WBS ²	NBS ³	SBS ⁴	EBS ⁵
188	<i>Acartia discaudata</i> (Giesbrecht, 1882)	+	+		+	
189	<i>Acartia longiremis</i> (Lilljeborg, 1853)	+	+	+	+	+
190	<i>Acartia tonsa</i> Dana, 1849	+	+	+	+	+
191	<i>Calanus finmarchicus</i> (Gunner, 1765)	+	+		+	+
192	<i>Calanus hyperboreus</i> Krøyer, 1838		+			
193	<i>Candacia armata</i> (Boeck, 1872)		+			
194	<i>Centropages chierchiae</i> Giesbrecht, 1889		+			
195	<i>Centropages hamatus</i> (Lilljeborg, 1853)	+	+	+	+	+
196	<i>Centropages typicus</i> Krøyer, 1849		+		+	
197	<i>Diaptomus graciloides</i> Lilljeborg, 1888 °				+	+
198	<i>Eudiaptomus gracilis</i> (G.O. Sars, 1862)		+		+	+
199	<i>Eurytemora affinis</i> (Poppe, 1880)	+	+	+	+	+
200	<i>Eurytemora carolleae</i> Alekseev and Souissi, 2011 [Recently described as a sibling species of <i>E. affinis</i> in the Baltic Sea; in the earlier studies in the BP, WBS and NBS, might have been confused with its sibling (Alekseev et al. 2009; Sukhikh et al., 2013)]				+	+
201	<i>Eurytemora hirundoides</i> (Nordquist, 1888) (Syn.: <i>Eurytemora hirundo</i> Giesbrecht, 1881)	+	+	+	+	+
202	<i>Eurytemora lacustris</i> (Poppe, 1887)				+	+
203	<i>Eurytemora velox</i> (Lilljeborg, 1853)		+			+
204	<i>Hetercope appendiculata</i> Sars, 1863 °					+
205	<i>Limnocalanus grimaldii</i> (De Guerne, 1886)	+	+	+	+	+
206	<i>Limnocalanus macrurus</i> G.O. Sars, 1863	+	+	+	+	+
207	<i>Metridia lucens</i> Boeck, 1865		+			
208	<i>Microcalanus pusillus</i> G.O. Sars, 1903		+			
209	<i>Paracalanus parvus</i> (Claus, 1863)	+	+		+	+
210	<i>Paraeuchaeta norvegica</i> (Boeck, 1872)	+	+			
211	<i>Pareucalanus attenuatus</i> (Dana, 1849)		+			
212	<i>Pseudocalanus acuspis</i> (Giesbrecht, 1881) [The recent molecular data (RFLP results) confirm that <i>P. acuspis</i> is the only <i>Pseudocalanus</i> species with a resident population in the Baltic Sea (Holmborn et al., 2011). Earlier, two more species from this genus were also considered for this sea: <i>Pseudocalanus elongatus</i> (Boeck, 1865) and <i>P. minutus</i> (Krøyer, 1845)]	+	+	+	+	+
213	<i>Temora longicornis</i> (O.F. Müller, 1785)	+	+	+	+	+

No	Taxa	BP ¹	WBS ²	NBS ³	SBS ⁴	EBS ⁵
Copepoda, Cyclopoida						
214	<i>Acanthocyclops robustus</i> (G.O. Sars, 1863)		+			
215	<i>Cyclopina gracilis</i> Claus, 1863		+			
216	<i>Cyclopina kieferi</i> Schäfer, 1936 **		+			
217	<i>Cyclopina norvegica</i> Boeck, 1864		+			
218	<i>Cyclops</i> sp.	+		+	+	+
219	<i>Cyclops americanus</i> Marsh, 1893					+
220	<i>Cyclops furcifer</i> Claus, 1857 °					+
221	<i>Cyclops kolensis</i> Lilljeborg, 1901 °					+
222	<i>Cyclops scutifer</i> G.O. Sars, 1863 °				+	+
223	<i>Cyclops strenuus</i> Fischer, 1851		+		+	+
224	<i>Cyclops vicinus</i> Ulyanin, 1875		+		+	+
225	<i>Diacyclops bicuspidatus</i> (Claus, 1857)		+		+	+
226	<i>Diacyclops bisetosus</i> (Rehberg, 1880)		+		+	+
227	<i>Ectocyclops phaleratus</i> (Koch, 1838) °					+
228	<i>Eucyclops graciloides</i> Lilljeborg, 1888		+			
229	<i>Eucyclops lilljeborgi</i> Sars, 1918 °					+
230	<i>Eucyclops macruroides</i> (Lilljeborg, 1901) °				+	+
231	<i>Eucyclops macrurus</i> (G.O. Sars, 1863)				+	+
232	<i>Eucyclops serrulatus</i> (Fischer, 1851)		+		+	+
233	<i>Eucyclops speratus</i> (Lilljeborg, 1901)		+			
234	<i>Halicyclops affinis</i> (G.O. Sars, 1863) **		+			
235	<i>Halicyclops magniceps</i> (Lilljeborg, 1853)		+		+	
236	<i>Halicyclops neglectus</i> Kiefer, 1935		+			
237	<i>Macrocyclus albidus</i> (Jurine, 1820)				+	+
238	<i>Macrocyclus fuscus</i> (Jurine, 1820) °				+	
239	<i>Megacyclops viridis</i> (Jurine, 1820)		+	+	+	+
240	<i>Megacyclops gigas</i> (Claus, 1857) °				+	
241	<i>Mesocyclops</i> sp.	+				
242	<i>Mesocyclops dybowskii</i> (Lande, 1890) °				+	
243	<i>Mesocyclops hyalinus</i> (Rehberg, 1880) **		+			
244	<i>Mesocyclops leuckarti</i> (Claus, 1857)		+	+	+	+
245	<i>Microcyclus bicolor</i> (Sars) **, °					+
246	<i>Microcyclus gracilis</i> (Lilljeborg) **, °					+
247	<i>Microcyclus varicans</i> (G. O. Sars, 1863) °					+
248	<i>Oithona atlantica</i> Farran, 1908		+		+	

No	Taxa	BP ¹	WBS ²	NBS ³	SBS ⁴	EBS ⁵
249	<i>Oithona similis</i> Claus, 1866	+	+		+	+
250	<i>Paracyclops</i> sp.	+				
251	<i>Paracyclops affinis</i> (G.O. Sars, 1863) [◦]					+
252	<i>Paracyclops fimbriatus</i> (Fischer, 1853) [◦]				+	+
253	<i>Thermocyclops crassus</i> (Fischer, 1853) [◦]				+	+
254	<i>Thermocyclops oithonoides</i> Sars, 1863		+	+	+	+
255	<i>Tropocyclops prasinus</i> (Fischer, 1860) [◦]				+	
Copepoda, Monstrilloida						
256	<i>Cymbasoma rigidum</i> Thompson, 1888 **		+			
257	<i>Cymbasoma thompsoni</i> (Giesbrecht, 1892)**		+			
258	<i>Monstrilla helgolandica</i> (Claus, 1863)		+			
Copepoda, Harpacticoida						
259	<i>Canthocamptus staphylinus</i> (Jurine, 1820)		+		+	+
260	<i>Dactilopodia vulgaris</i> (Sars) **, [◦]					+
261	<i>Ectinosoma melaniceps</i> Boeck, 1865		+			+
262	<i>Halectinosoma curticorne</i> (Boeck, 1872)	+	+		+	
263	Harpacticoida indet.	+	+	+	+	+
264	<i>Microsetella norvegica</i> (Boeck, 1865)		+		+	+
265	<i>Microarthridion littorale</i> (Pope, 1881) [◦]	+				
266	<i>Nitokra hibernica</i> (Brady, 1880) [◦]				+	
267	<i>Onychocamptus mohammed</i> (Blanchard and Richard, 1891) (Syn.: <i>Laophonte mohammed</i> Blanchard and Richard, 1891) [◦]				+	
Pteropoda						
268	<i>Limacina retroversa</i> (Fleming, 1823)	+	+			
Polychaeta						
269	<i>Bylgides sarsi</i> (Kinberg in Malmgren, 1865)	+	+	+		+
270	<i>Harmothoe imbricata</i> (Linnaeus, 1769)		+			
271	<i>Harmothoe impar</i> (Johnston, 1839)		+			+
272	<i>Nephtys</i> sp.		+			+
273	<i>Nereis diversicolor</i> O.F. Müller, 1776		+	+		
274	<i>Pygospio elegans</i> Claparède, 1863	+	+	+		+
275	<i>Tomopteris helgolandica</i>	+	+			
	Polychaeta, larvae	+	+	+	+	+

No	Taxa	BP ¹	WBS ²	NBS ³	SBS ⁴	EBS ⁵
Chaetognatha						
276	<i>Parasagitta elegans</i> (Verrill, 1873)	+	+		+	+
277	<i>Parasagitta setosa</i> (Mueller, 1847)	+	+			+
278	<i>Sagitta bipunctata</i> Quoy & Gaimard, 1828		+			
Copelata						
279	<i>Fritillaria borealis</i> Lohmann, 1896	+	+	+	+	+
280	<i>Oikopleura dioica</i> Fol, 1872	+	+		+	+
	Larvae of Bivalvia (Mollusca) (Plate 5.3.31)	+	+	+	+	+
	Larvae of Gastropoda (Mollusca) (Plate 5.3.31)	+	+	+	+	+
	Larvae of Cirripedia (Crustacea) (Plate 5.3.32)	+	+	+	+	+
	Larvae of Bryozoa (Plate 5.3.35)	+	+	+	+	+
	Larvae of Echinodermata (Plate 5.3.35)	+	+	+	+	

¹ **BP:** **Baltic Proper:** after Ackefors (1965, 1969), Ostenfeld (1931), Mankowski (1948b, 1950b, 1951, 1959), Siudzinski (1965), Mielck & Künne (1932-1935); Holmborn et al., 2011;

² **WBS:** **Western Baltic Sea** (Kieler Bight, Mecklenburg Bight): after Remane (1940), Gerlach (2000), Kube et al. (2007a, b); Holmborn et al., 2011;

³ **NBS:** **Northern Baltic Sea** (Archipelago Sea, Bothnian Sea): after Vuorinen (pers. com.), Lindquist (1959), Ostenfeld (1931); Holmborn et al., 2011;

⁴ **SBS:** **Southern Baltic Sea** (Darss-Zingst Bodden Chain; Gdansk Basin; Vistula Lagoon; Curonian Lagoon): after Mankowski (1948a, 1950a, b), Telesh & Heerkloss (2002, 2004, and references therein); Alekseev et al., 2009; Holmborn et al., 2011; Sukhikh et al., 2013;

⁵ **EBS:** **Eastern Baltic Sea** (Gulf of Riga; Gulf of Finland, *including* the freshwater Neva Bay): after Purasjoki (1958), Flinkman (pers. com.), Telesh (pers. com.), Silina (1997), Telesh & Heerkloss (2002, 2004, and references therein), Rodionova et al. (2005), Rodionova & Panov (2006); Alekseev et al., 2009; Holmborn et al., 2011; Sukhikh et al., 2013.

*Syn. – synonyms; ** – not registered in ITIS; ^c – species found mainly in the coastal nearshore (oligohaline or fresh) waters